

BICROSSED PRODUCTS FOR FINITE GROUPS

A. L. AGORE, A. CHIRVĂȘITU, B. ION, AND G. MILITARU

Dedicated to Freddy Van Oystaeyen on the occasion of his 60th birthday.

ABSTRACT. We investigate one question regarding bicrossed products of finite groups which we believe has the potential of being approachable for other classes of algebraic objects (algebras, Hopf algebras). The problem is to classify the groups that can be written as bicrossed products between groups of fixed isomorphism types. The groups obtained as bicrossed products of two finite cyclic groups, one being of prime order, are described.

INTRODUCTION

The bicrossed product construction is a generalization of the semidirect product construction for the case when neither factor is required to be normal: a group E is the internal bicrossed product of its subgroups H and G if $HG = E$ and their intersection is trivial. Groups with this property (but allowing for nontrivial intersection) have been in the literature for a quite long time under the terminology *permutable groups* [7, 8] or groups that admit an *exact factorization* (see e.g. [3, 14]).

The bicrossed product construction itself is due to Zappa [13]. It was rediscovered by Szép [11] and yet again by Takeuchi [12]. The terminology *bicrossed product* is taken from Takeuchi, other terms referring to this construction used in the literature are *knit product* and *Zappa-Szép product*. Bicrossed product constructions were subsequently introduced and studied for other structures: algebras, Hopf algebras, Lie algebras, Lie groups, locally compact quantum groups, groupoids. For Hopf algebras, in particular, structural results are still missing and objects obtained from such constructions form a considerable proportion of the known examples (see e.g. [1]). Assume for simplicity that k is a field of characteristic zero. Let E be a finite group that is a bicrossed product of the groups H and G . A noncommutative noncocommutative Hopf algebra $k[H]^* \# k[G]$ that is both semisimple and cosemisimple can be constructed [12]. This is the easiest way to construct semisimple cosemisimple finite dimensional Hopf algebras. For this reason we decided to investigate some aspects of the bicrossed product construction in its original finite group setting.

Our main question, going back to Ore [8] asks for the description of all groups which arise as bicrossed products of two fixed groups. Little progress has been made on this question. In this respect we would like to mention the result of Wielandt [15] establishing that from

1991 *Mathematics Subject Classification*. 20B05, 20B35, 20D06, 20D40.

The work of A. L. Agore, B. Ion and G. Militaru where partially supported by CNCSIS grant 24/28.09.07 of PN II "Groups, quantum groups, corings and representation theory". The work of B. Ion was partially supported by NSF grant DMS-0536962.

two finite nilpotent groups of coprime orders one always obtains a solvable group and the work of Douglas [2] on finite groups expressible as bicrossed products of two finite cyclic groups. Finding all matched pairs between two finite cyclic groups seems to be still an open question, even though J. Douglas [2] has devoted four papers and over two dozen theorems to the subject. In fact, solving this problem does not provide an answer to the classification of all associated bicrossed products and does not indicate whether a bicrossed product could not be obtained more easily as a semidirect product. In Section 2 we will give a complete answer to this question for the case of two finite cyclic groups, one of them being of prime order. As it turns out, if a group is isomorphic to a bicrossed product of two finite cyclic groups, one of them being of prime order then it is isomorphic to a semidirect product between the same cyclic groups. We would like also to mention some interesting recent investigations [4], [6] into the corresponding question at the level of algebras.

1. PRELIMARIES

1.1. Definitions and notation. Let us fix the notation that will be used throughout the paper. Let H and G be two groups and $\alpha : G \times H \rightarrow H$ and $\beta : G \times H \rightarrow G$ two maps. We use the notation

$$\alpha(g, h) = g \triangleright h \quad \text{and} \quad \beta(g, h) = g \triangleleft h$$

for all $g \in G$ and $h \in H$. The map α (resp. β) is called trivial if $g \triangleright h = h$ (resp. $g \triangleleft h = g$) for all $g \in G$ and $h \in H$. If $\alpha : G \times H \rightarrow H$ is an action of G on H as group automorphisms we denote by $H \rtimes_{\alpha} G$ the semidirect product of H and G : $H \rtimes_{\alpha} G = H \times G$ as a set with the multiplication given by

$$(h_1, g_1) \cdot (h_2, g_2) := (h_1(g_1 \triangleright h_2), g_1 g_2)$$

for all $h_1, h_2 \in H, g_1, g_2 \in G$.

The opposite group structure on H will be denoted by H^{op} : $H^{\text{op}} = H$ as a set with the multiplication $h_1 \cdot_{\text{op}} h_2 = h_2 h_1$ for all $h_1, h_2 \in H$.

Definition 1.1. A *matched pair* of groups is a quadruple $\Lambda = (H, G, \alpha, \beta)$ where H and G are groups, $\alpha : G \times H \rightarrow H$ is a left action of the group G on the set H , $\beta : G \times H \rightarrow G$ is a right action of the group H on the set G such that the following compatibility conditions hold:

$$g \triangleright (h_1 h_2) = (g \triangleright h_1)((g \triangleleft h_1) \triangleright h_2) \tag{1}$$

$$(g_1 g_2) \triangleleft h = (g_1 \triangleleft (g_2 \triangleright h))(g_2 \triangleleft h) \tag{2}$$

for all $g, g_1, g_2 \in G, h, h_1, h_2 \in H$.

A morphism $\varphi : (H_1, G_1, \alpha_1, \beta_1) \rightarrow (H_2, G_2, \alpha_2, \beta_2)$ between two matched pairs consists of a pair of group morphisms $\varphi_H : H_1 \rightarrow H_2, \varphi_G : G_1 \rightarrow G_2$ such that

$$\varphi_H \circ \alpha_1 = \alpha_2 \circ (\varphi_G \times \varphi_H), \quad \varphi_G \circ \beta_1 = \beta_2 \circ (\varphi_G \times \varphi_H)$$

Remark 1.2. Let $\Lambda = (H, G, \alpha, \beta)$ be a matched pair of groups. Then

$$g \triangleright 1 = 1 \quad \text{and} \quad 1 \triangleleft h = 1 \tag{3}$$

for all $g \in G$ and $h \in H$.

Let H and G be groups and $\alpha : G \times H \rightarrow H$ and $\beta : G \times H \rightarrow G$ two maps. Let $H \rtimes_{\alpha \bowtie \beta} G = H \rtimes G := H \times G$ as a set with an binary operation defined by the formula:

$$(h_1, g_1) \cdot (h_2, g_2) := (h_1(g_1 \triangleright h_2), (g_1 \triangleleft h_2)g_2) \quad (4)$$

for all $h_1, h_2 \in H, g_1, g_2 \in G$.

The main motivation behind the definition of matched pair is the following result (we refer to [12] or [5, section IX.1] for the proof).

Theorem 1.3. *Let H and G be groups and α and β two maps as above. Then $H \rtimes_{\alpha \bowtie \beta} G$ is a group with unit $(1, 1)$ if and only if (H, G, α, β) is a matched pair. Moreover, a morphism between two matched pairs induces a morphism between the corresponding groups.*

If (H, G, α, β) is a matched pair the group $H \rtimes G$ is called the *bicrossed product* (or the *Zappa-Szép product*) of H and G . The inverse of an element of the group $H \rtimes G$ is given by the formula

$$(h, g)^{-1} = (g^{-1} \triangleright h^{-1}, (g \triangleleft (g^{-1} \triangleright h^{-1}))^{-1}) \quad (5)$$

for all $h \in H$ and $g \in G$. Also, remark that $H \times \{1\} \cong H$ and $\{1\} \times G \cong G$ are subgroups of $H \rtimes G$ and every element (h, g) of $H \rtimes G$ can be written uniquely as a product of an element of $H \times \{1\}$ and of an element of $\{1\} \times G$ as follows:

$$(h, g) = (h, 1) \cdot (1, g) \quad (6)$$

Conversely, one can see that this observation characterizes the bicrossed product. Again, we refer to [5, 12] for the details.

Theorem 1.4. *Let E be a group $H, G \leq E$ be subgroups such that any element of E can be written uniquely as a product of an element of H and an element of G . Then there exists a matched pair (H, G, α, β) such that*

$$\theta : H \rtimes G \rightarrow E, \quad \theta(h, g) = hg$$

is group isomorphism.

The maps α and β play in fact a symmetric role.

Proposition 1.5. *Let $\Lambda = (H, G, \alpha, \beta)$ be a matched pair of groups. Then*

- (i) $\tilde{\Lambda} = (G, H, \tilde{\alpha}, \tilde{\beta})$, where $\tilde{\alpha}$ and $\tilde{\beta}$ are given by

$$\tilde{\alpha} : H \times G \rightarrow G, \quad \tilde{\alpha}(h, g) = \left(\beta(g^{-1}, h^{-1}) \right)^{-1} \quad (7)$$

$$\tilde{\beta} : H \times G \rightarrow H, \quad \tilde{\beta}(h, g) = \left(\alpha(g^{-1}, h^{-1}) \right)^{-1} \quad (8)$$

for all $h \in H$ and $g \in G$ is a matched pair of groups.

- (ii) *The map*

$$\chi : \left(H \rtimes_{\alpha \bowtie \beta} G \right)^{\text{op}} \rightarrow G \rtimes_{\tilde{\alpha} \bowtie \tilde{\beta}} H, \quad \chi(h, g) = (g^{-1}, h^{-1})$$

is a group isomorphism. In particular,

$$\xi : H \rtimes_{\alpha \bowtie \beta} G \rightarrow G \rtimes_{\tilde{\alpha} \bowtie \tilde{\beta}} H, \quad \xi(h, g) = \left(g \triangleleft (g^{-1} \triangleright h^{-1}), (g^{-1} \triangleright h^{-1})^{-1} \right)$$

is a group isomorphism.

Proof. The proof is a straightforward verification. $\square$

Remark 1.6. Let H and G be two groups as above and let $\beta : G \times H \rightarrow G$ be the trivial action. Then (H, G, α, β) is a matched pair if and only if the map $\alpha : G \times H \rightarrow H$ is an action of G on H as group automorphisms. In this case the bicrossed product is the semidirect product $H \rtimes_{\alpha} G$.

Assume now that the map α is the trivial action. We obtain from (8) that $\tilde{\beta}$ is trivial. Keeping in mind that the bicrossed product $G \rtimes_{\tilde{\alpha}} \rtimes_{\tilde{\beta}} H$ with the trivial $\tilde{\beta}$ is a semidirect product we can invoke Proposition 1.5 (ii) to conclude that $H \rtimes G \cong G \rtimes_{\tilde{\alpha}} H$.

1.2. Universality properties. Let $\Lambda = (H, G, \alpha, \beta)$ be a matched pair of groups. We associate to Λ two categories such that the bicrossed product of H and G becomes an initial object in one of them and a final object in the other.

Define the category ${}_{\Lambda}\mathcal{C}$ as follows: the objects of ${}_{\Lambda}\mathcal{C}$ are pairs $(X, (u, v))$ where X is a group, $u : H \rightarrow X$, $v : G \rightarrow X$ are group morphisms such that:

$$v(g)u(h) = u(g \triangleright h)v(g \triangleleft h) \quad (9)$$

for all $g \in G$, $h \in H$. A morphism in ${}_{\Lambda}\mathcal{C}$

$$f : (X, (u, v)) \rightarrow (X', (u', v'))$$

is a morphism of groups $f : X \rightarrow X'$ such that $f \circ u = u'$ and $f \circ v = v'$. It can be checked that $(H \rtimes G, (i_H, i_G))$ is an object in ${}_{\Lambda}\mathcal{C}$, where i_H and i_G are the canonical inclusions of H and G inside their bicrossed product.

Define the category $\mathcal{C}_{\Lambda}$ as follows: the objects of $\mathcal{C}_{\Lambda}$ are pairs $(X, (u, v))$ where X is a group, $u : X \rightarrow H$, $v : X \rightarrow G$ are two maps such that the following two compatibility condition holds:

$$u(xy) = u(x)(v(x) \triangleright u(y)), \quad v(xy) = (v(x) \triangleleft u(y))v(y) \quad (10)$$

for all $x, y \in X$. A morphism in $\mathcal{C}_{\Lambda}$

$$f : (X, (u, v)) \rightarrow (X', (u', v'))$$

is a morphism of groups $f : X \rightarrow X'$ such that $u' \circ f = u$ and $v' \circ f = v$. It can be checked that $(H \rtimes G, (\pi_H, \pi_G))$ is an object in $\mathcal{C}_{\Lambda}$, where p_H and p_G are the canonical projections from the bicrossed product to H and G .

Proposition 1.7. Let $\Lambda = (H, G, \alpha, \beta)$ be a matched pair of groups. Then

- (i) $(H \rtimes G, (i_H, i_G))$ is an initial object of ${}_{\Lambda}\mathcal{C}$.
- (ii) $(H \rtimes G, (\pi_H, \pi_G))$ is a final object of $\mathcal{C}_{\Lambda}$.

Proof. (i) Let $(X, (u, v)) \in {}_{\Lambda}\mathcal{C}$. We have to prove that there exists a unique morphism of groups $w : H \rtimes G \rightarrow X$ such that $w \circ i_H = u$ and $w \circ i_G = v$.

Assume that w satisfies this condition. Then using (6) we have:

$$\begin{aligned} w((h, g)) &= w((h, 1) \cdot (1, g)) = w((h, 1))w((1, g)) \\ &= (w \circ i_H)(h)(w \circ i_G)(g) = u(h)v(g) \end{aligned}$$

for all $h \in H$ and $g \in G$ and this proves that w is unique.

If we define

$$w : H \bowtie G \rightarrow X, \quad w(h, g) = u(h)v(g)$$

then

$$\begin{aligned} w((h_1, g_1) \cdot (h_2, g_2)) &= w(h_1(g_1 \triangleright h_2), (g_1 \triangleleft h_2)g_2) \\ &= u(h_1)u(g_1 \triangleright h_2)v(g_1 \triangleleft h_2)v(g_2) \\ &= u(h_1)v(g_1)u(h_2)v(g_2) \\ &= w((h_1, g_1))w((h_2, g_2)) \end{aligned}$$

showing that w is a morphism of groups.

Part (ii) follows by a similar argument. $\square$

Straightforward from Proposition 1.7 we obtain the description of morphisms between a group and a bicrossed product.

Corollary 1.8. *Let E be a group and (H, G, α, β) a matched pair. Then*

- (i) *$w : H \bowtie G \rightarrow E$ is a group morphism if and only if there exist $u : H \rightarrow E$ and $v : G \rightarrow E$ group morphisms such that*

$$v(g)u(h) = u(g \triangleright h)v(g \triangleleft h), \quad \text{and} \quad w(h, g) = u(h)v(g)$$

for all $h \in H$ and $g \in G$.

- (ii) *$w : E \rightarrow H \bowtie G$ is a morphism of groups if and only if there exist $u : E \rightarrow H$ and $v : E \rightarrow G$ two maps such that*

$$u(xy) = u(x)(v(x) \triangleright u(y)), \quad v(xy) = (v(x) \triangleleft u(y))v(y)$$

and $w(x) = (u(x), v(x))$ for all $x, y \in E$.

Remark 1.9. Corollary 1.8 can be used to describe all morphisms or isomorphisms between two matched pairs $H \alpha \bowtie \beta G$ and $H \alpha' \bowtie \beta' G$. However, the descriptions are rather technical and we will not include them here.

2. BICROSSED PRODUCTS BETWEEN FINITE CYCLIC GROUPS

As mentioned in the Introduction the question of describing all groups which arise as bicrossed products of two given groups was asked by Ore. The first and, by our knowledge, the only systematic study of this kind, for groups which arise as bicrossed products of two finite cyclic groups, was employed by J. Douglas in 1951. In his first paper on the subject [2, pag. 604] Douglas formulates the problem he wants to solve: describe all groups all whose elements are expressible in the form $a^i b^j$ where a and b are independent elements of order n and, respectively, m . What Douglas refers to as independent elements is in fact the condition that the cyclic groups generated by each of these elements have trivial intersection. Therefore the problem can be formulated as follows: describe all groups which arise as bicrossed products of two finite cyclic groups.

In what follows C_n and C_m will be two cyclic groups of orders n and m . We denote by a and b a fixed generator of C_n and, respectively, C_m . For any positive integer k we denote

by $\mathbb{Z}_k$ the ring of residue classes modulo k and by $S(\mathbb{Z}_k)$ the set of bijective functions from $\mathbb{Z}_k$ to itself. Let $\alpha : C_m \times C_n \rightarrow C_n$, and $\beta : C_m \times C_n \rightarrow C_m$ be two actions. They are completely determined by two maps $\theta \in S(\mathbb{Z}_n)$ and $\phi \in \mathbb{Z}_m$ such that

$$\alpha(b, a^x) = a^{\theta(x)}, \quad \beta(b^y, a) = b^{\phi(y)}$$

for any $x \in \mathbb{Z}_n$, $y \in \mathbb{Z}_m$. Douglas [2, Theorem I] obtained necessary and sufficient conditions on the pair of maps $(\theta, \phi) \in S(\mathbb{Z}_n) \times S(\mathbb{Z}_m)$ which would make $(C_n, C_m, \alpha, \beta)$ a matched pair. The functions θ and ϕ satisfying his conditions were called *conjugate special substitutions*.

Finding all pairs of conjugate special substitutions (or, equivalently, all matched pairs between two finite cyclic groups) seems to be still an open question. Furthermore, solving this problem does not provide an answer to the classification problem for the associated bicrossed products. In particular, it does not indicate whether a bicrossed product could not be obtained more easily as a semidirect product.

We investigate the structure of bicrossed products of two finite cyclic groups, one of which has prime order. We shall prove now our main result.

Theorem 2.1. *Let p be a prime number, and $m \geq 2$ a positive integer. A bicrossed product between two cyclic groups of orders p and respectively m is isomorphic to a semidirect product between cyclic groups of the same orders p and m .*

Proof. Let G be a group, C_p and C_m two fixed cyclic subgroups of G , of orders p and m respectively, such that $G = C_p \rtimes C_m$. Let a, b be generators for C_p and C_m respectively. We can certainly assume that C_p and C_m are not normal in G .

Let $H = C_m \cap aC_ma^{-1}$. A subgroup of a finite cyclic group is uniquely determined by its order so $aHa^{-1} = H$. Since C_m and aC_ma^{-1} are abelian, H is central in both. Also note that $C_m \neq aC_ma^{-1}$, because we assumed C_m is not normal in G . The centralizer $C(H)$ of H in G must then strictly contain C_m . Since C_m has index p in G , it is maximal. It follows that $C(H) = G$, that is H is a central subgroup of G .

In fact, for any $1 \leq k \leq p-1$, H is the intersection between C_m and $a^k C_m a^{-k}$. Indeed, a^k generates C_p , so any subgroup of G normalized by a^k must also be normalized by a .

We are now going to work in the quotient group G/H . Let us denote by

$$\pi : G \rightarrow G/H$$

the canonical projection. For any $1 \leq k \leq p-1$, the groups $\pi(C_m)$ and $\pi(a^k C_m a^{-k})$ intersect trivially. We can now apply a theorem of Frobenius [10, Theorem 9.11 and Exercise 9.9] for the group G/H to conclude that the subgroup $\pi(C_p)$ is normal. This means that $\pi(bab^{-1}) = \pi(a^t)$ for some $1 \leq t \leq p-1$, or, equivalently, that

$$bab^{-1} = a^t c$$

for some $c \in H$. Raising this identity to power p , we find that $c^p = 1$. We may assume that $c \neq 1$, otherwise C_p would be normal in G . Also, $t \neq 1$ otherwise $a^{-1}ba = cb \in C_m$, which means that C_m is normal in G , contrary to our assumption. Hence, we can find an integer u such that $u(t-1) \equiv 1 \pmod{p}$.

With the notation $\tilde{a} = ac^u$, we then have $b\tilde{a}b^{-1} = \tilde{a}^t$. The proof is now finished: the subgroup of G generated by $\tilde{a}$ is normal in G , has order p and intersects C_m trivially. $\square$

ACKNOWLEDGMENTS: We thank the referee for detailed suggestions that have helped us improve this paper.

REFERENCES

- [1] S. Caenepeel, B. Ion, G. Militaru, and S. Zhu, The factorization problem and the smash biproduct of algebras and coalgebras. *Algebr. Represent. Theory* **3** (2000), no. 1, 19–42.
- [2] J. Douglas, On finite groups with two independent generators. I, II, III, IV. *Proc. Nat. Acad. Sci. U. S. A.* **37** (1951), 604–610, 677–691, 749–760, 808–813.
- [3] M. Giudici, Factorisations of sporadic simple groups. *J. Algebra* **304** (2006), no. 1, 311–323.
- [4] P. Jara, J. López Peña, G. Navarro, D. Stefan, On the classification of twisting maps between K^n and K^m , arXiv:0805.2874.
- [5] C. Kassel, Quantum groups. *Graduate Texts in Mathematics* **155**. Springer-Verlag, New York, 1995.
- [6] J. López Peña and G. Navarro, On the classification and properties of noncommutative duplicates. *K-Theory* **38** (2008), no. 2, 223–234.
- [7] E. Maillet, Sur les groupes échangeables et les groupes décomposables. *Bull. Soc. Math. France* **28** (1900), 7–16.
- [8] O. Ore, Structures and group theory. I. *Duke Math. J.* **3** (1937), no. 2, 149–174.
- [9] L. Rédei, Zur Theorie der faktorisierbaren Gruppen, I. *Acta Math. Acad. Sci. Hungar.* **1** (1950), 74–98.
- [10] J. Rotman, An introduction to the theory of groups. Fourth edition. *Graduate Texts in Mathematics* **148**. Springer-Verlag, New York, 1995.
- [11] J. Szép, Über die als Produkt zweier Untergruppen darstellbaren endlichen Gruppen. *Comment. Math. Helv.* **22** (1949), 31–33.
- [12] M. Takeuchi, Matched pairs of groups and bismash products of Hopf algebras. *Comm. Algebra* **9** (1981), no. 8, 841–882.
- [13] G. Zappa, Sulla costruzione dei gruppi prodotto di due dati sottogruppi permutabili tra loro. *Atti Secondo Congresso Un. Mat. Ital., Bologna, 1940*, 119–125. Edizioni Cremonense, Rome, 1942.
- [14] J. Wiegold and A. G. Williamson, The factorisation of the alternating and symmetric groups. *Math. Z.* **175** (1980), no. 2, 171–179.
- [15] H. Wielandt, Über das Produkt paarweise vertauschbarer nilpotenter Gruppen. *Math. Z.* **55** (1951) 1–7.

(A.L.A., A.C., G.M.) FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF BUCHAREST, STR. ACADEMIEI 14, RO-70109 BUCHAREST 1, ROMANIA

E-mail address: ana.agore@fmi.unibuc.ro, chirvasitua@gmail.com, gigel.militaru@fmi.unibuc.ro

(B.I.) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PITTSBURGH, PITTSBURGH, PA 15260